

Phase diagram of disordered fermion model on two-dimensional square lattice with π -flux

Takahiro Fukui*

Department of Mathematical Sciences, Ibaraki University, Mito 310-8512, Japan

(February 1, 2008)

A fermion model with random on-site potential defined on a two-dimensional square lattice with π -flux is studied. The continuum limit of the model near the zero energy yields Dirac fermions with random potentials specified by four independent coupling constants. The basic symmetry of the model is time-reversal invariance. Moreover, it turns out that the model has enhanced (chiral) symmetry on several surfaces in the four-dimensional space of the coupling constants. It is shown that one of the surfaces with chiral symmetry has $\text{Sp}(n) \times \text{Sp}(n)$ symmetry whereas others have $\text{U}(2n)$ symmetry, both of which are broken to $\text{Sp}(n)$, and the fluctuation around a saddle point is described, respectively, by $\text{Sp}(n)_2$ WZW model and $\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model. Based on these results, we propose a phase diagram of the model.

PACS: 72.15.Rn, 71.23.-k, 11.10.-z

I. INTRODUCTION

Anderson localization has attracted renewed interest since a plenty of new universality classes were discovered [1–3]. One is a system with particle-hole symmetry studied by Gade [1]. Although generic disordered systems in two dimensions are believed to be insulators [4], she showed that such a system has a random critical point at the band center. Typical examples studied so far are the random flux model [5–8] and the random-hopping model with π -flux [9,10,7,8]. This class is often referred to as *AIII* and *BDI* in the case with broken and unbroken time reversal symmetry, respectively, which corresponds to the chiral Gaussian unitary and chiral Gaussian orthogonal ensemble of the random matrix theory in the zero dimensional limit [2].

Other examples of new universality classes are found in disordered d -wave superconductors described by a mean-field Bogoliubov-deGennes Hamiltonian [11–23]. Altland and Zirnbauer [3,2] classified them by two basic symmetries, i.e, spin-rotation and time-reversal into four kinds of classes. With (without) spin-rotation invariance, disordered d -wave superconductors belong to class *C* and *CI* (*D* and *DIII*), respectively, in the case with broken and unbroken time-reversal symmetry. The beta function of the renormalization group for corresponding nonlinear sigma models in two dimensions tells that the classes *C* and *CI* have localized states only, whereas the classes *D* and *DIII* may have delocalized states. What is particularly interesting is that some of them allow in the nonlinear sigma model description the topological term [15,16,23] which plays a crucial role, for example, in quantum Hall transitions [24], or the Wess-Zumino-Witten (WZW) term [17,19,23] which gives rise to a nontrivial strong-coupling fixed point. Therefore, disordered d -wave superconductors are expected to have quite rich phase diagram.

These developments can shed new light on the localization problems. For example, the question whether the density of states of the random flux model diverges had been a long-standing problem. Recognizing, however, that it belongs to the Gade's class of *AIII* type, Furusaki concluded that the DOS actually diverges. Another example is the disordered fermion model with π -flux we will revisit in this paper. This model was already studied by Fisher and Fradkin [25] in 1984. They showed that the effective field theory of the lattice model with random on-site potentials belongs to the universality class of $\text{O}(2n, 2n)/\text{O}(2n) \times \text{O}(2n)$ nonlinear sigma model, concluding thereby that all states are localized.

A point here is that the continuum limit of the model includes random potentials specified by four independent coupling constants. Since Fisher and Fradkin studied only the case where all the coupling constants are equal, we can expect richer phase diagram in the whole space of the coupling constants. Actually, although the generic points in this space should flow into the localized phase mentioned-above, the model has enhanced symmetry in special cases and accordingly renormalization group flows in them are toward different fixed points. Another point in this paper is concerned with chiral symmetry. It was shown that generic disordered d -wave superconductors in class *CI*, described by Dirac fermions, yield no WZW term [17,23] although those Dirac fermions have chiral $\text{Sp}(n) \times \text{Sp}(n)$ symmetry. The reason may be that chirality is not unique in the d -wave Dirac fermions, since the lattice fermion gives the species doubling in the continuum limit. Here, chiral symmetry means that there exist a matrix which anticommutes with a Hamiltonian.

In this paper, we investigate in detail the phase diagram of the disordered fermion model with π -flux by means of the fermionic replica method to clarify these points. We show that this model, described by double Dirac fermions, has three kinds of chiral symmetry in special cases. Interestingly, these different chiral symmetries lead to different universality classes. To be concrete, an effective action of the Green functions has $\text{Sp}(n) \times \text{Sp}(n)$ symmetry in one of them and $\text{U}(2n)$ symmetry in the others, both of which are broken down to $\text{Sp}(n)$. It turns out that corresponding Goldstone mode can be described by $\text{Sp}(n)_2$ WZW model and $\text{U}(n)/\text{Sp}(n)$ nonlinear sigma model, respectively.

This paper is organized as follows. In the next section, we introduce a lattice model, which is the same as Fisher and Fradkin studied, and derive Dirac Hamiltonian in the continuum limit. Although the most generic model has only time-reversal invariance, symmetry is enhanced on three surfaces in the space of the coupling constants. We focus our attention to one of them with $\text{Sp}(n) \times \text{Sp}(n)$ symmetry (others have $\text{U}(2n)$ symmetry) and give calculations in detail. Based on the continuum Hamiltonian, we derive an effective generating functional of ensemble-averaged Green functions in Sec. III. We show that $\text{Sp}(n) \times \text{Sp}(n)$ symmetry is spontaneously broken down to $\text{Sp}(n)$. In Sec. IV, we derive an effective action for the Goldstone mode, resulting in $\text{Sp}(n)_2$ WZW model ($\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model for the others). In Sec. V, we present the phase diagram. In Sec. VI, we give a summary and concluding remarks.

II. MODEL

A. A lattice model and its continuum limit

We define a basic lattice model on the square lattice in two dimensions. For convenience, we introduce similar notations used by Ludwig et al [26] in the following way. The square lattice in two dimensions is divided into unit




FIG. 1. The square lattice on which the model is defined. Unit cells (plaquettes) are enclosed by the dotted-lines. Four sites on each plaquette are numbered counterclockwise. Thick lines means that nearest neighbor hopping is negative on them due to π flux.

cells, each composed of a plaquette enclosing four sites. A plaquette is labeled by a set of integers $j \equiv (j_x, j_y)$ and four sites on it are numbered 1, 2, 3 and 4 counterclockwise, as illustrated in Fig. 1. With such a notational convention, a pure and disorder Hamiltonian denoted respectively as H_p and H_d are defined by

$$\begin{aligned}
 H_p &= \sum_j \left(c_{1j}^\dagger c_{2j} + c_{2j}^\dagger c_{3j} - c_{3j}^\dagger c_{4j} + c_{4j}^\dagger c_{1j} + c_{2j}^\dagger c_{1j+\hat{x}} - c_{3j}^\dagger c_{4j+\hat{x}} + c_{3j}^\dagger c_{2j+\hat{y}} + c_{4j}^\dagger c_{1j+\hat{y}} + \text{h.c.} \right), \\
 H_d &= \sum_j \sum_{a=1}^4 v'_{aj} c_{aj}^\dagger c_{aj},
 \end{aligned} \tag{2.1}$$

where $\hat{x} \equiv (1, 0)$, $\hat{y} \equiv (0, 1)$, and v'_{aj} is random real number due to nonmagnetic impurities. Negative signs of the third and the sixth terms in H_p are due to π flux imposed. Now let us construct an effective Hamiltonian near the

band center in the case of weak disorder. To this end, we first derive the continuum limit of the pure model, and next we take into account the effects of disorder.

Fourier transformation $c_{aj} = \int \frac{d^2k}{(2\pi)^2} e^{ikj} c_a(k)$ for $a = 1, \dots, 4$ leads to

$$H_p = \int \frac{d^2k}{(2\pi)^2} c^\dagger(k) \begin{pmatrix} 1 + e^{-ik_x} & 1 + e^{-ik_y} & 1 + e^{-ik_x} \\ 1 + e^{ik_x} & 1 + e^{ik_y} & -(1 + e^{ik_x}) \\ 1 + e^{ik_y} & -(1 + e^{-ik_x}) & \end{pmatrix} c(k),$$

where $c^\dagger(k) \equiv (c_1^\dagger(k), \dots, c_4^\dagger(k))$. It is readily seen that the Hamiltonian gives zero-energies at $k = (\pi, \pi)$. In the vicinity of them, the model can be effectively described by Dirac fermions. To see this, set $k = (\pi, \pi) + a_0 p$ with the lattice constant a_0 , expand the Hamiltonian to the first order of p , and introduce a field operator in the continuum limit $c_a((\pi, \pi) + a_0 p) \equiv a_0^{-\frac{3}{2}} \psi_a(p)$ for $a = 1, \dots, 4$. Then we have

$$H_p \sim \int \frac{d^2p}{(2\pi)^2} \psi^\dagger(p) (-\sigma_2 \otimes 1 p_x - \sigma_1 \otimes \sigma_2 p_y) \psi(p),$$

where $\psi^\dagger = (\psi_1^\dagger, \psi_2^\dagger, \psi_3^\dagger, \psi_4^\dagger)$. Accordingly, effective Hamiltonian near the zero energy can be expressed by the following field theoretical Hamiltonian

$$H_p = \int d^2x \psi^\dagger(x) \mathcal{H}_p \psi(x), \quad \mathcal{H}_p \equiv \gamma_\mu i \partial_\mu, \quad (2.2)$$

where $x = a_0 j$, $\psi(x) \equiv \int \frac{d^2p}{(2\pi)^2} e^{ipx} \psi(p)$, and γ matrices are defined as $\gamma_1 = \sigma_2 \otimes 1$ and $\gamma_2 = \sigma_1 \otimes \sigma_2$.

By using the relation between the lattice operator c_{aj} and the continuum field operator $\psi_a(x)$,

$$c_{aj} \sim \int \frac{d^2(a_0 p)}{(2\pi)^2} e^{i(\pi, \pi)j + ia_0 p j} a_0^{-\frac{3}{2}} \psi_a(p) = a_0^{\frac{1}{2}} (-)^{\frac{x}{a_0} + \frac{y}{a_0}} \psi_a(x),$$

the continuum limit of the disorder potentials are readily derived; $H_d = \int d^2x \sum_{a=1}^4 \psi_a^\dagger(x) v'_a(x) \psi_a(x)$, where $v'_a(x)$ is the slowly-varying part of v'_{aj} defined as $v'_{aj} \equiv a_0 v'_a(x)$. Taking suitable linear combination of v'_a , we can rewrite the random potentials as

$$H_d = \int d^2x \psi^\dagger(x) \mathcal{H}_d \psi(x), \quad \mathcal{H}_d = \sum_{\nu=1}^4 v_\nu(x) \alpha_\nu, \quad (2.3)$$

where $\alpha_1 = 1 \otimes \sigma_3, \alpha_2 = \sigma_3 \otimes \sigma_3, \alpha_3 = \sigma_3 \otimes 1$ and $\alpha_4 = 1 \otimes 1$. We assume that the four disorder potentials obey Gaussian probability distribution

$$P[v_\nu(x)] \propto e^{-\frac{2}{g_\nu} v_\nu^2(x)}, \quad \text{for } \nu = 1, \dots, 4. \quad (2.4)$$

It may be more convenient to switch into another basis via suitable rotation for ψ ,

$$\begin{aligned} \gamma_1 &= \sigma_2 \otimes 1, & \gamma_2 &= \sigma_1 \otimes \sigma_3, \\ \alpha_1 &= 1 \otimes \sigma_2, & \alpha_2 &= \sigma_3 \otimes \sigma_2, & \alpha_3 &= \sigma_3 \otimes 1, & \alpha_4 &= 1 \otimes 1. \end{aligned} \quad (2.5)$$

In what follows, we refer to the space where these matrices live as V . It is distinguished from the replica space W_r introduced momentarily. The total Hamiltonian we investigate in this paper is $H = H_p + H_d$ in Eqs. (2.2) and (2.3) with (2.5). It is noted that this is the same model as Fisher and Fradkin [25] studied in the case $g_1 = g_2 = g_3 = g_4$. They showed that the effective field theory is $O(2n, 2n)/O(2n) \times O(2n)$ nonlinear sigma model for the bosonic replicas [or $Sp(2n)/Sp(n) \times Sp(n)$ nonlinear sigma model for the fermionic ones], concluding thereby that all states are localized. According to Zirnbauer, this class is referred to as AII [2]. Although a slight relaxation of the condition $g_1 = g_2 = g_3 = g_4$ still leads to the same universality class, several subspaces in (g_1, g_2, g_3, g_4) have not only time-reversal symmetry but also additional enhanced symmetry specified below, and therefore belong to different universality classes.

B. Symmetries of the model

In this subsection, we examine symmetries of the present model, which plays a crucial role in the classification of the universality classes for disordered systems. First of all, the most general Hamiltonian has unbroken time-reversal symmetry. In the original basis, it is expressed by $\mathcal{H}^t = \mathcal{H}$, whereas in the rotated basis where the Hamiltonian is described by the matrices in Eq. (2.5), it is expressed by

$$\mathcal{H} = \mathcal{T}\mathcal{H}^t\mathcal{T}^{-1}, \quad \mathcal{T} = 1 \otimes \sigma_1. \quad (2.6)$$

If the model does not have any other symmetries, the universality class is *AII* and hence all states are localized. There are several subspaces in (g_1, g_2, g_3, g_4) , however, on which symmetry is enhanced. To be concrete, the Hamiltonian (2.2) has following symmetry

$$\mathcal{H} = -\mathcal{P}_j\mathcal{H}\mathcal{P}_j^{-1}, \quad \text{with} \quad \begin{cases} \mathcal{P}_1 = \sigma_3 \otimes \sigma_3, & \text{for } g_3 = g_4 = 0 \\ \mathcal{P}_2 = \sigma_1 \otimes \sigma_1, & \text{for } g_2 = g_4 = 0 \\ \mathcal{P}_3 = \sigma_1 \otimes i\sigma_2, & \text{for } g_1 = g_4 = 0 \end{cases}. \quad (2.7)$$

Recently, Hatsugai et.al [9] proposed the same model as Eq. (2.1) but with random hoppings instead of random on-site potential, which we will refer to as HWKM model. They have shown that such Hamiltonian has not only time-reversal symmetry but also another one often called chiral symmetry. It has been shown [9,10,7,8] that this symmetry is responsible for the existence of delocalized critical point at the zero energy. Chiral symmetry in this case is expressed as $\mathcal{H} = -\mathcal{P}_4\mathcal{H}\mathcal{P}_4^{-1}$ with $\mathcal{P}_4 = \sigma_3 \otimes 1$, similarly to Eq. (2.7), and the universality class is specified as *BDI* or in other words $U(2n)/Sp(n)$ ($Gl(2n, R)/O(2n)$) nonlinear sigma model for fermionic (bosonic) replica method. Considering these, it is easily guessed that the symmetry (2.7) also plays a crucial role in Anderson localization for the present model. In what follows, we refer to the symmetry in Eq. (2.7) also as chiral symmetry.

III. REPLICA METHOD

It is so tedious to repeat similar calculations that we concentrate on the case $\mathcal{P}_1$, and hence we often denote $\mathcal{P}_1$ just as $\mathcal{P}$ for short. But with slight modification, any $\mathcal{P}_j$ can apply to the formulation. At the moment, the difference between $\mathcal{P}_1$ and the others is whether $\mathcal{P}_j$ commutes or anticommutes with $\mathcal{T}$ and whether it is real symmetric or antisymmetric. Surprisingly, this difference, which seems trivial at first glance, leads to a different universality class.

In order to study the transport properties of the model near the zero energy, we need to compute the Green functions

$$G(\pm i\epsilon) = \frac{1}{\pm i\epsilon - \mathcal{H}}.$$

With the $\mathcal{P}$ symmetry, however, we do not need to distinguish the retarded and advanced Green functions, since they are related with each other via the relation [27],

$$G(+i\epsilon) = \mathcal{P} \frac{1}{i\epsilon + \mathcal{H}} \mathcal{P}^{-1} = -\mathcal{P}G(-i\epsilon)\mathcal{P}^{-1}.$$

The Green function $G(i\epsilon)$ can be computed by a replicated functional

$$\mathcal{Z} = \int \mathcal{D}\psi \mathcal{D}\psi^\dagger e^{-\int d^2x \mathcal{L}},$$

with

$$\mathcal{L} = \sum_{\alpha=1}^n \sum_{j,k=1}^4 \psi_{j\alpha}^* (i\epsilon 1_4 - \mathcal{H})_{jk} \psi_{k\alpha} \equiv \text{tr}_{W_r} \psi^\dagger (i\epsilon 1_4 - \mathcal{H}) \psi, \quad (3.1)$$

where the subscript α denotes the n species in the replica space W_r , $\psi_{\alpha j}^\dagger = \psi_{j\alpha}^*$, and the usual convention of the matrix product is applied to the fermion matrix field $\psi_{j\alpha}^*$ and $\psi_{j\alpha}$. Although such a notational convention is not always necessary, it is quite convenient for us to save many complicated subscripts. It should be noted that the fields $\psi_{\alpha j}^*$ and $\psi_{\alpha j}$ are completely independent Grassmann variables. Therefore, they obey in general different transformation laws.

A. Auxiliary space

Based on the Lagrangian (3.1), we will construct an effective theory of order parameter fields for the replicas by averaging-over disorder. In this process, it is quite important to fully incorporate the symmetries (2.6) and (2.7) in the space V . To this end, we introduce an auxiliary space to reflect these into the replicas, according to the method developed by Zirnbauer [2]. Namely, we extend the replica space as $W_r \rightarrow W \equiv W_r \otimes W_t \otimes W_c$, where W_t (W_c) stands for the space associated with time-reversal (chiral) symmetry. We then introduce a $4n \times 4$ ($4 \times 4n$) matrix field $\tilde{\Psi}_{\alpha j}$ ($\Psi_{j\alpha}$),

$$\tilde{\Psi} = \begin{pmatrix} \tilde{\chi}_+ \\ \tilde{\chi}_- \end{pmatrix} \quad \Psi = (\chi_+, \chi_-); \quad \tilde{\chi}_{\pm} = \tilde{\chi} \mathcal{P}_{\mp}, \quad \chi_{\pm} = \mathcal{P}_{\pm} \chi, \quad (3.2)$$

where

$$\tilde{\chi} = \frac{1}{\sqrt{2}} \begin{pmatrix} \psi^\dagger \\ -\psi^t \mathcal{T} \end{pmatrix}, \quad \chi = \frac{1}{\sqrt{2}} (\psi, \mathcal{T} \psi^*), \quad (3.3)$$

$\mathcal{P}_{\pm} = \frac{1 \pm \mathcal{P}}{2}$, and ψ^t stands for the transpose of ψ . Since the components of these fields are not independent of each other, the fields are subject to the constraints

$$\begin{aligned} \tilde{\Psi} &= \gamma \Psi^t \mathcal{T}, \quad \Psi = -\mathcal{T} \tilde{\Psi}^t \gamma^{-1}, \quad \gamma = 1_n \otimes i\sigma_2 \otimes 1_2, \\ \tilde{\Psi} &= -\pi \tilde{\Psi} \mathcal{P}, \quad \Psi = \mathcal{P} \Psi \pi, \quad \pi = 1_n \otimes 1_2 \otimes \sigma_3. \end{aligned} \quad (3.4)$$

Note that the matrices γ and π are defined in W , whereas $\mathcal{T}$ and $\mathcal{P}$ are defined in V . We immediately find

$$\Psi \tilde{\Psi} = \mathcal{T} (\Psi \tilde{\Psi})^t \mathcal{T}^{-1}, \quad \Psi \tilde{\Psi} = -\mathcal{P} (\Psi \tilde{\Psi}) \mathcal{P}^{-1}. \quad (3.5)$$

These relations tell that the matrix $\Psi \tilde{\Psi}$ in V has the same symmetries as $\mathcal{H}$ has. It is also easy to find alternative relations for the matrix $\tilde{\Psi} \Psi$ in W ,

$$\tilde{\Psi} \Psi = \gamma (\tilde{\Psi} \Psi)^t \gamma^{-1}, \quad \tilde{\Psi} \Psi = -\pi (\tilde{\Psi} \Psi) \pi^{-1}. \quad (3.6)$$

These relations actually specify the symmetry of the model, as we shall see later.

Although the potentials α_1 and α_2 have time-reversal and chiral symmetries, the converse is not true: The symmetries (2.6) and (2.7) admit other potentials $\beta_1 = \sigma_1 \otimes 1$, $\beta_2 = \sigma_2 \otimes \sigma_3$, $\beta_3 = 1 \otimes \sigma_1$ and $\beta_4 = \sigma_3 \otimes \sigma_1$. The reason why there are no beta matrices in Eq. (2.3) is that the basic lattice model (2.1) includes only on-site random potential. Namely, if we take all kinds of possible disorder for the lattice Hamiltonian including random (nearest neighbor and diagonal) hoppings as well as random on-site potentials, we have a continuum Hamiltonian with all kinds of possible disorder potentials specified by the relations (2.6) and (2.7). Such a model may be referred to as a model with maximum entropy. Since we are interested in the model with on-site random potentials, we have to remove these β matrices. For this purpose, we take account of additional conditions

$$\mathcal{H}_d = -\mathcal{T} \mathcal{H}_d \mathcal{T}, \quad \mathcal{H}_d = \mathcal{E} \mathcal{H}_d \mathcal{E}, \quad \mathcal{E} \equiv \sigma_3 \otimes 1. \quad (3.7)$$

It is readily seen that the potentials including beta matrices do not satisfy these relations. Let us then introduce the fields

$$\tilde{\Psi}^\circ = \frac{1}{2} \begin{pmatrix} \tilde{\Psi} \\ \tilde{\Psi} \mathcal{E} \\ -\tilde{\Psi} \mathcal{T} \\ -\tilde{\Psi} \mathcal{E} \mathcal{T} \end{pmatrix}, \quad \Psi^\circ = \frac{1}{2} (\Psi, \mathcal{E} \Psi, \mathcal{T} \Psi, \mathcal{T} \mathcal{E} \Psi). \quad (3.8)$$

We refer to this extra space as W° . It is easy to verify that the matrix $\Psi^\circ \tilde{\Psi}^\circ$ obeys the same relations that $\mathcal{H}_d$ does in (3.7) and in (3.5) with extension $\gamma \rightarrow \gamma \otimes 1_2 \otimes \sigma_3$ and $\pi \rightarrow \pi \otimes 1_2 \otimes \sigma_3$, which means that the undesirable potentials are in fact excluded. By using these fields we can reexpress the Lagrangian (3.1) as

$$\mathcal{L} = \text{tr}_W (i\epsilon \omega \tilde{\Psi} \Psi - \tilde{\Psi} i \not{\partial} \Psi) - \text{tr}_{W \otimes W^\circ} \tilde{\Psi}^\circ \mathcal{H}_d \Psi^\circ, \quad (3.9)$$

where $\omega \equiv 1_n \otimes 1_2 \otimes \sigma_1$ and $\not{\partial} \equiv \gamma_\mu \partial_\mu$.

B. Effective Lagrangian averaged-over disorder

Now we are ready to ensemble-average the Lagrangian (3.9) over disorder. In this subsection, we assume $g_1 = g_2 = g$ for simplicity. More general case $g_1 \neq g_2$ is briefly discussed in Sec. IIID. The probability distribution (2.4) can be expressed as

$$P[\mathcal{H}_d] \propto e^{-\frac{1}{2g} \text{tr}_V \mathcal{H}_d^2}. \quad (3.10)$$

Therefore, the Lagrangian is converted into

$$\begin{aligned} \mathcal{L} &= \frac{1}{2g} \text{tr}_V \mathcal{H}_d^2 + \text{tr}_W (i\epsilon\omega \tilde{\Psi}\Psi - \tilde{\Psi}i\partial\Psi) - \text{tr}_{W \otimes W^\circ} \tilde{\Psi}^\circ \mathcal{H}_d \Psi^\circ, \\ &= \frac{1}{2g} \text{tr}_V \left(\mathcal{H}_d + g\Psi^\circ \tilde{\Psi}^\circ \right)^2 - \frac{g}{2} \text{tr}_V (\Psi^\circ \tilde{\Psi}^\circ)^2 + \text{tr}_W (i\epsilon\omega \tilde{\Psi}\Psi - \tilde{\Psi}i\partial\Psi), \end{aligned}$$

where we have used a relation $\text{tr}_{W \otimes W^\circ} \tilde{\Psi}^\circ \mathcal{H}_d \Psi^\circ = -\text{tr}_V \mathcal{H}_d \Psi^\circ \tilde{\Psi}^\circ$. It should be noted that integration over $\mathcal{H}_d$ is readily performed, since the matrix $\Psi^\circ \tilde{\Psi}^\circ$ has completely the same symmetries as $\mathcal{H}_d$ has. This is the reason why we have introduced several auxiliary spaces in the last subsection. By using $\text{tr}_V (\Psi^\circ \tilde{\Psi}^\circ)^2 = -\text{tr}_{W \otimes W^\circ} (\tilde{\Psi}^\circ \Psi^\circ)^2$ again and the definition of (3.8), the gaussian integration over $\mathcal{H}_d$ leads to

$$\mathcal{L} = \text{tr}_W (i\epsilon\omega \tilde{\Psi}\Psi - \tilde{\Psi}i\partial\Psi) + \frac{g'}{2} \sum_{\tau=1}^4 \text{tr}_W (\tilde{\Psi} \mathcal{O}_\tau \Psi)^2, \quad (3.11)$$

where $\mathcal{O}_1 = 1, \mathcal{O}_2 = \mathcal{E}, \mathcal{O}_3 = i\mathcal{T}, \mathcal{O}_4 = i\mathcal{T}\mathcal{E}$, and $g' = \frac{g}{4}$. We mention in passing that if the model have maximum entropy, there appears only one interaction term with $\mathcal{O}_1 = 1$ and $g' = g$. Eq. (3.11) is an effective Lagrangian which generates the ensemble-averaged Green functions for the present model. It is easy to confirm that new matrices are subject to

$$\tilde{\Psi} \mathcal{O}_\tau \Psi = \gamma (\tilde{\Psi} \mathcal{O}_\tau \Psi)^t \gamma^{-1}, \quad \tilde{\Psi} \mathcal{O}_\tau \Psi = \mp \pi (\tilde{\Psi} \mathcal{O}_\tau \Psi) \pi^{-1} \quad \text{for} \quad \begin{cases} \tau = 1, 2 \\ \tau = 3, 4, \end{cases} \quad (3.12)$$

So far we have obtained an effective action including four fermi interactions by averaging over disorder. The next task is to introduce auxiliary (order-parameter) fields to decouple them. To this end, we add the following terms of auxiliary fields Q_τ ($\tau = 1, \dots, 4$) to the Lagrangian (3.11):

$$\mathcal{L}' = -\frac{1}{2g'} \sum_{\tau=1}^4 \text{tr}_W (Q_\tau + g' \tilde{\Psi} \mathcal{O}_\tau \Psi + \delta_{\tau 1} i\epsilon\omega)^2. \quad (3.13)$$

Then, the Lagrangian becomes

$$\mathcal{L} = -\text{tr}_W \tilde{\Psi} i\partial\Psi - \frac{1}{2g'} \sum_{\tau} \text{tr}_W (Q_\tau^2 + i\epsilon\omega Q_1 \delta_{\tau 1}) - \sum_{\tau} \text{tr}_W Q_\tau \tilde{\Psi} \mathcal{O}_\tau \Psi. \quad (3.14)$$

Eq. (3.13) requires that Q_τ should obey the same symmetries that $\tilde{\Psi} \mathcal{O}_\tau \Psi$ does in Eq. (3.12). Therefore,

$$Q_\tau^\dagger = -Q_\tau, \quad Q_\tau = \gamma Q_\tau^t \gamma^{-1}, \quad Q_\tau = \mp \pi Q_\tau \pi^{-1} \quad \text{for} \quad \begin{cases} \tau = 1, 2 \\ \tau = 3, 4, \end{cases} \quad (3.15)$$

The antihermiticity ensures that the integration over Q_τ in Eq. (3.14) converges.

This Lagrangian (3.14) has a global $\text{Sp}(n) \times \text{Sp}(n)$ symmetry. To see this, let us consider a transformation $\Psi \rightarrow \Psi g^{-1}$, $\tilde{\Psi} \rightarrow g \tilde{\Psi}$, and $Q_\tau \rightarrow g Q_\tau g^{-1}$. Since this transformation should keep the relations (3.15), g is subject to $g^\dagger g = 1, g^t \gamma g = \gamma$ and $g^{-1} \pi g = \pi$. Therefore, g has the following form

$$g = \begin{pmatrix} g_+ & \\ & g_- \end{pmatrix}, \quad g_\pm g_\pm^\dagger = 1, \quad g_\pm \gamma_0 g_\pm^t = \gamma_0, \quad \gamma_0 \equiv 1_n \otimes i\sigma_2,$$

where $2n \times 2n$ matrices $g_\pm$ belongs to $W_{\text{rt}} = W_{\text{r}} \otimes W_{\text{t}}$ and the explicit matrix above represent the space W_{c} . The conditions for $g_\pm$ require $g_\pm \in \text{Sp}(n)$.

Next let us solve Eq. (3.15),

$$Q_\tau = \begin{pmatrix} & M_\tau \\ -M_\tau^\dagger & \end{pmatrix}, \quad \text{for } \tau = 1, 2,$$

$$Q_\tau = \begin{pmatrix} A_{\tau-2}^+ & \\ & A_{\tau-2}^- \end{pmatrix}, \quad \text{for } \tau = 3, 4,$$

where $2n \times 2n$ matrices M_τ and $A_\tau^\pm$ are subject to $M_\tau = -\gamma_0 M_\tau^* \gamma_0^{-1}$, $A_\tau^\pm = \gamma_0 A_\tau^{\pm\dagger} \gamma_0^{-1}$, and $A_\tau^{\pm\dagger} = -A_\tau^\pm$. Using the definition in Eq. (3.2), the Lagrangian is now written as

$$\begin{aligned} \mathcal{L} = & -\text{tr}_{W_{\text{rt}}}(\tilde{\chi}_+ i \not{\partial} \chi_+ + \tilde{\chi}_- i \not{\partial} \chi_-) + \frac{1}{2g'} \sum_{\tau=1}^2 \text{tr}_{W_{\text{rt}}} [2M_\tau M_\tau^\dagger - (A_\tau^{+2} + A_\tau^{-2})] - \frac{i\epsilon}{2g'} \text{tr}_{W_{\text{rt}}}(M_1 - M_1^\dagger) \\ & - \sum_{\tau=1}^2 \text{tr}_{W_{\text{rt}}} (M_\tau \tilde{\chi}_- \mathcal{O}_\tau \chi_+ - M_\tau^\dagger \tilde{\chi}_+ \mathcal{O}_\tau \chi_- + A_\tau^+ \tilde{\chi}_+ \mathcal{O}_{\tau+2} \chi_+ + A_\tau^- \tilde{\chi}_- \mathcal{O}_{\tau+2} \chi_-). \end{aligned} \quad (3.16)$$

The tranformation laws under g are

$$\chi_\pm \rightarrow \chi_\pm g_\pm^\dagger, \quad \tilde{\chi}_\pm \rightarrow g_\pm \tilde{\chi}_\pm, \quad M_\tau \rightarrow g_+ M_\tau g_-^\dagger, \quad A_\tau^\pm \rightarrow g_\pm A_\tau^\pm g_\pm^\dagger. \quad (3.17)$$

So far we have examined the case $\mathcal{P}_1$. It has been shown that the Lagrangian has $\text{Sp}(n) \times \text{Sp}(n)$ symmetry and hence the model belongs to CI , which is the same class as d -wave superconductors with unbroken time-reversal and spin-rotation symmetries. Contrary to this, in the other cases associated with $\mathcal{P}_2$ and $\mathcal{P}_3$, symmetry of the Lagrangian is quite different. Similar calculations above show that these Lagrangians have $U(2n)$ symmetry insted of $\text{Sp}(n) \times \text{Sp}(n)$ dut to the fact that γ defined in (3.4) is $\gamma = 1_n \otimes i\sigma_2 \otimes \sigma_1$ if we take the same π . This is the same class as HKWM model, which is often referred to as BDI or chiral orthogonal class.

C. Effective potential

Based on the Lagrangian in the last subsection, we derive the effective potential to study the spontaneous symmetry breaking. We assume that the order-paramters are diagonal in the space W_{r} . Namely, set

$$M_\tau = i \text{diag}(m_{\tau 1}, \dots, m_{\tau n}) \otimes 1_2, \quad A_\tau^\pm = i \text{diag}(a_{\tau 1} \pm b_{\tau 1}, \dots, a_{\tau n} \pm b_{\tau n}) \otimes 1_2, \quad \tau = 1, 2,$$

where $m_{\tau\alpha}(x)$, $a_{\tau\alpha}(x)$ and $b_{\tau\alpha}(x)$ are real fields. Then the Lagrangian (3.16) becomes

$$\mathcal{L} = \frac{2}{g'} \sum_{\tau=1}^2 \sum_{\alpha=1}^n (m_{\tau\alpha}^2 + a_{\tau\alpha}^2 + b_{\tau\alpha}^2) + \frac{2\epsilon}{g'} \sum_{\alpha=1}^n m_{1\alpha} + \sum_{\alpha=1}^n \psi_\alpha^\dagger (-i \not{\partial} + v_\alpha) \psi_\alpha,$$

where v_α is 4×4 matrix defined by

$$v_\alpha = -i(m_{1\alpha} 1 \otimes 1 + m_{2\alpha} \sigma_3 \otimes 1) + a_{1\alpha} 1 \otimes \sigma_1 - ib_{1\alpha} \sigma_3 \otimes \sigma_2 + a_{2\alpha} \sigma_3 \otimes \sigma_1 - ib_{1\alpha} 1 \otimes \sigma_2.$$

Integrating out the fermi fields, we have an effective potential $V_\alpha(\{m, a, b\}) = S(\{m(x) = m, a(x) = a, b(x) = b\})/\text{volume of one-loop order}$,

$$V_\alpha(\{m, a, b\}) = \frac{2}{g'} \sum_{\tau=1}^2 \sum_{\alpha=1}^n (m_{\tau\alpha}^2 + a_{\tau\alpha}^2 + b_{\tau\alpha}^2) - \int \frac{d^2 k}{(2\pi)^2} \ln \det_V(\not{k} + v_\alpha) + \frac{2\epsilon}{g'} \sum_{\alpha=1}^n m_{1\alpha}.$$

Since the potential V_α is a function of six variables, it is hard to find out the true minimum. However, it is readily expected from the symmetry breaking term that $m_{1\alpha}$ has a finite value. Assuming it, we have tried to find out nontrivial sultion $\partial V_\alpha / \partial m_{\tau\alpha} = \partial V_\alpha / \partial a_{\tau\alpha} = \partial V_\alpha / \partial b_{\tau\alpha} = 0$, but in vain. Although possibility of some other solutions still remains, the solution $m_{1\alpha} \neq 0$ and others=0 is of course a natural solution. In this case, the saddle point equaiton for $m_{1\alpha}$ becomes

$$0 = \frac{\partial V_\alpha}{\partial m_{1\alpha}} = \frac{4m_{1\alpha}}{g'} - 2 \int \frac{d^2 k}{(2\pi)^2} \frac{2m_{1\alpha}}{k^2 + m_{1\alpha}^2},$$

which gives nontrivial solution $m_{1\alpha} = m \equiv \Lambda(e^{\frac{4\pi}{g'}} - 1)^{\frac{1}{2}} \sim \Lambda e^{-\frac{2\pi}{g'}}$. This result suggests that the operator $(\tilde{\Psi}\mathcal{O}_1\Psi)^2$ is relevant whereas others are irrelevant around the present saddle-point. Therefore, in the following analysis, we neglect the irrelevant interactions and take only the relevant interaction into account. In the next section, we will discuss slow fluctuation around the saddle point. But before it, we will see how the anisotropy $g_1 \neq g_2$ modifies the theory so far obtained.

D. Anisotropy effects

In the case where $g_1 = g_2$, we could treat them simultaneously in Eq. (3.10). Contrary to this simple case, we have to treat the two potentials α_1 and α_2 separately in anisotropic case. To this end, first introduce a new matrix $\mathcal{F} = \sigma_1 \otimes 1_2$ under which they transform

$$\alpha_1 = \mathcal{F}\alpha_1\mathcal{F}^{-1}, \quad \alpha_2 = -\mathcal{F}\alpha_2\mathcal{F}^{-1}.$$

We next introduce

$$\tilde{\Psi}_1^\circ = \frac{1}{\sqrt{2}} \begin{pmatrix} \tilde{\Psi}^\circ \\ \tilde{\Psi}^\circ \mathcal{F} \end{pmatrix}, \quad \tilde{\Psi}_2^\circ = \frac{1}{\sqrt{2}} \begin{pmatrix} \tilde{\Psi}^\circ \\ -\tilde{\Psi}^\circ \mathcal{F} \end{pmatrix}, \quad \Psi_1^\circ = \Psi_2^\circ = \frac{1}{\sqrt{2}}(\Psi^\circ, \mathcal{F}\Psi^\circ).$$

Let us denote $\mathcal{H}_{dj} \equiv v_j \alpha_j$ for $j = 1, 2$ and the extra auxiliary space as W' . Since the matrix $\Psi_j^\circ \tilde{\Psi}_j^\circ$ ($j = 1, 2$) has completely the same symmetry property as $\mathcal{H}_{dj}$, we can average over disorder in the following way:

$$\begin{aligned} & \frac{1}{2g_1} \text{tr}_V \mathcal{H}_{d1}^2 + \frac{1}{2g_2} \text{tr}_V \mathcal{H}_{d2}^2 - \text{tr}_{W \otimes W^\circ \otimes W'} \left(\tilde{\Psi}_1^\circ \mathcal{H}_{d1} \Psi_1^\circ + \tilde{\Psi}_2^\circ \mathcal{H}_{d2} \Psi_2^\circ \right) \\ & \rightarrow \frac{g_1}{2} \text{tr}_{W \otimes W^\circ \otimes W'} (\tilde{\Psi}_1^\circ \Psi_1^\circ)^2 + \frac{g_2}{2} \text{tr}_{W \otimes W^\circ \otimes W'} (\tilde{\Psi}_1^\circ \Psi_1^\circ)^2 \\ & = \frac{g_1 + g_2}{2} \text{tr}_{W \otimes W^\circ} (\tilde{\Psi}^\circ \Psi^\circ)^2 + \frac{g_1 - g_2}{2} \text{tr}_{W \otimes W^\circ} (\tilde{\Psi}^\circ \mathcal{F} \Psi^\circ)^2, \end{aligned}$$

where arrow means that the integration over $\mathcal{H}_{dj}$ is carried out. This equation shows that anisotropy $g_1 \neq g_2$ yields an additional term including $\mathcal{F}$. However, it turns out that this term is also irrelevant and therefore can be neglected, following similar discussions to the previous subsection.

IV. EFFECTIVE ACTION FOR THE GOLDSTONE MODE

The saddle-point $M_1 = m 1_n \otimes 1_2$ is invariant under $g_+ = g_-$ type transformation in Eq. (3.17) but not invariant under $g_+ = g_-^\dagger$. This means that $\text{Sp}(n) \times \text{Sp}(n)$ symmetry is broken to $\text{Sp}(n)$. Therefore, fluctuation around the saddle-point can be incorporated by parametrizing the order-parameter field M_1 as

$$M_1 = \xi i m \xi = i m \xi^2 = i m U,$$

where ξ and $U \in \text{Sp}(n)$, i.e., $U^t \gamma_0 U = \gamma_0$, $U^\dagger U = 1_{2n}$. The field U describes the Goldstone mode around the saddle point. Then, the Lagrangian becomes

$$\mathcal{L} = -\text{tr}_{W_{\text{rt}}} [\tilde{\chi}_+ i \not{\partial} \chi_+ + \tilde{\chi}_- i \not{\partial} \chi_- + i m (U \tilde{\chi}_- \chi_+ + U^\dagger \tilde{\chi}_+ \chi_-)] + \frac{\epsilon m}{2g'} \text{tr}_{W_{\text{rt}}} (U + U^\dagger) + \frac{2nm^2}{g'}$$

The last constant term will be neglected, since it vanishes in the replica limit $n \rightarrow 0$. In the following subsection, we construct an effective theory for small fluctuation U around the saddle point.

A. Integration over fermi fields

So far we have treated $\tilde{\chi}$ and χ as matrix fields to save complicated subscripts. It has actually worked well especially for the discussions of the symmetry. However, in order to extract an effective Lagrangian for U via chiral gauge transformation introduced below, we cannot treat the space V and W separately any longer. Therefore, we

switch into the standard notation for fermi fields as follows: Note the following identity, for example, $\text{tr}_{W_{\text{rt}}} U \tilde{\chi}_- \chi_+ = \sum_{\alpha, \alpha'} \sum_j U_{\alpha' \alpha} \tilde{\chi}_{-\alpha j} \chi_{+j \alpha'} = \sum_{j, \alpha} \sum_{j', \alpha'} \tilde{\chi}_{-\alpha j} \delta_{jj'} \otimes U_{1\alpha\alpha'}^\dagger \chi_{+j' \alpha'} = \sum_{j, \alpha} \sum_{j', \alpha'} \tilde{\chi}_{-\alpha j} \delta_{jj'} \otimes \gamma_0 U_{1\alpha\alpha'}^\dagger \gamma_0^{-1} \chi_{+j' \alpha'} \rightarrow \tilde{\chi}_- 1_4 \otimes U^\dagger \chi_+$, where the last arrow means that we have made a γ_0 rotation to $\tilde{\chi}$ and χ and we have regarded $\chi_\pm$ as a vector with $8n$ components in $V \otimes W_{\text{rt}}$. Then the fermion part of the Lagrangian can be denoted as

$$\begin{aligned} -\mathcal{L}_F &= \tilde{\chi}_+ i \not{\partial} \otimes 1_{2n} \chi_+ + \tilde{\chi}_- i \not{\partial} \otimes 1_{2n} \chi_- + im (\tilde{\chi}_- 1_4 \otimes U^\dagger \chi_+ + \tilde{\chi}_+ 1_4 \otimes U \chi_-) \\ &= \tilde{\chi} [i \not{\partial} \otimes 1_{2n} + im (\mathcal{P}_+ \otimes U^\dagger + \mathcal{P}_- \otimes U)] \chi. \end{aligned} \quad (4.1)$$

Although this Lagrangian seems to be standard, the integration over χ is not straightforward. It is because the elementary fields $\psi^\dagger$ and ψ are doubly included in $\tilde{\chi}$ and χ defined in Eq. (3.3). It is possible, however, to rewrite this Lagrangian as

$$-\mathcal{L}_F = \tilde{\psi}' \begin{pmatrix} imU^\dagger & i(\partial_2 - i\partial_1)1_{2n} \\ i(\partial_2 + i\partial_1)1_{2n} & imU \end{pmatrix} \psi'; \quad \tilde{\psi}' = (\psi_1^\dagger, -\psi_3^\dagger, \psi_2^\dagger, -\psi_4^\dagger), \quad \psi' = \begin{pmatrix} \psi_1 \\ \psi_3^* \\ \psi_2 \\ \psi_4^* \end{pmatrix}, \quad (4.2)$$

where $\tilde{\psi}'$ and ψ' have $4n$ components whose explicit subscripts in the above definition are those for V . It should be noted that this Lagrangian is described by independent field variables $\tilde{\psi}'$ and ψ' only. As a result, we can integrate out fermi fields, which results in

$$\begin{aligned} \int \mathcal{D}\psi \mathcal{D}\psi^\dagger e^{\int d^2x \mathcal{L}_F} &= \text{Det}_{V \otimes W_{\text{rt}}} \begin{pmatrix} imU^\dagger & i(\partial_2 - i\partial_1)1_{2n} \\ i(\partial_2 + i\partial_1)1_{2n} & imU \end{pmatrix} \\ &= \text{Det}_{V \otimes W_{\text{rt}}}^{\frac{1}{2}} [i \not{\partial} \otimes 1_{2n} + im (\mathcal{P}_+ \otimes U^\dagger + \mathcal{P}_- \otimes U)]. \end{aligned} \quad (4.3)$$

Based on this result, we derive an effective action for the fluctuation U in the next subsection.

B. Chiral gauge transformation

Suppose that the fluctuation $U(x)$ around the saddle-point varies slowly with x . Then, we can expand, with respect to derivatives, the determinant derived in the last subsection. To this end, make the folowing transformation

$$\begin{aligned} \tilde{\chi}_+^U &= \tilde{\chi}_+ U, & \chi_+^U &= U^\dagger \chi_+, \\ \tilde{\chi}_-^U &= \tilde{\chi}_-, & \chi_-^U &= \chi_-. \end{aligned} \quad (4.4)$$

Here and hereafter, we often denote $1_4 \otimes U$ just as U for simplicity. Without confusion, we also use e.g., $\mathcal{P}$ instead of $\mathcal{P} \otimes 1_{2n}$. The Lagrangian is now converted into

$$-\mathcal{L}_F = \tilde{\chi}^U (i \hat{\mathcal{D}} + im) \chi^U,$$

where

$$\hat{D}_\mu = D_{+\mu} + \partial_{-\mu}; \quad D_{+\mu} = D_\mu \mathcal{P}_+, \quad \partial_{-\mu} = \partial_\mu \mathcal{P}_-, \quad (4.5)$$

with

$$D_\mu = \partial_\mu + A_\mu, \quad A_\mu = U^\dagger \partial_\mu U.$$

Integration over fermi fields $\tilde{\chi}^U$ and χ^U gives also a determinant, and the action for U is

$$S = -\frac{1}{2} \ln \text{Det}_{V \otimes W_{\text{rt}}} (i \hat{\mathcal{D}} + im) - \ln J(U) + \frac{\epsilon m}{2g'} \int d^2x \text{tr}_{W_{\text{rt}}} (U + U^\dagger) \quad (4.6)$$

where $J(U)$ is a Jacobian [28] due to the transformation (4.4). It turns out that expansion of the determinant yields no second order terms with respect to the derivative, and hence we neglect the contribution from it in the leading order approximation. In what follows, we then concnetrate on the calculation of the Jacobian. We mention in passing that we will use the Lagrangian (4.1) to compute the Jacobian, although (4.2) is less confusing.

Let us consider the following eigenvalue problem for the Dirac operator (4.5) to define and regularize the Jacobian of the path integration,

$$i\hat{D}\varphi_n = \lambda_n\varphi_n, \quad \phi_n^\dagger i\hat{D} = \phi_n^\dagger \lambda_n,$$

where right- and left-eigenvalue equations are introduced, since the Dirac operator is non-hermitian. The eigenfunctions are normalized via the inner product $(\phi_n^\dagger, \varphi_m) \equiv \int d^2x \phi_n^\dagger \varphi_m = \delta_{nm}$. Assuming the completeness of the eigenfunctions, the fields $\tilde{\chi}$ and χ are expanded as

$$\begin{aligned} \chi^U &= \sum_n a_n^U \varphi_n, & a_n^U &= (\phi_n^\dagger, \chi^U), \\ \tilde{\chi}^U &= \sum_n \phi_n^\dagger \tilde{b}_n^U, & \tilde{b}_n^U &= (\tilde{\chi}^U, \varphi_n). \end{aligned}$$

The measure is then defined as

$$\mathcal{D}\chi^U = \prod_n da_n^U \quad \text{or} \quad \mathcal{D}\tilde{\chi}^U = \prod_n d\tilde{b}_n^U.$$

It should be noted that if one uses the Lagrangian (4.2) described by independent field variables $\tilde{\psi}'$ and ψ' only, the measure is defined by $\mathcal{D}\psi'\mathcal{D}\tilde{\psi}'$ as usual. In the present case, however, either $\mathcal{D}\chi$ or $\mathcal{D}\tilde{\chi}$ is enough, since each already includes all independent variables. It seems to be unusual, to be sure, but we follow the calculation with the latter notation. Using the definitions for the path integration measure, we can compute the Jacobian for infinitesimal small change for the gauge transformation (4.4), $\chi^U \rightarrow \chi^{U+\delta U} = \chi^U + \delta\chi^U$ and $\tilde{\chi}^U \rightarrow \tilde{\chi}^{U+\delta U} = \tilde{\chi}^U + \delta\tilde{\chi}^U$ with $\delta\chi^U = -\mathcal{P}_+ \otimes U^\dagger \delta U \chi^U$ and $\delta\tilde{\chi}^U = \tilde{\chi}^U \mathcal{P}_- \otimes U^\dagger \delta U$ as follows [28]:

$$\begin{aligned} a_n^{U+\delta U} &= (\phi_n^\dagger, \chi^U + \delta\chi^U) = \sum_m (1 - N)_{nm} a_m^U, \\ \tilde{b}_n^{U+\delta U} &= (\tilde{\chi}^U + \delta\tilde{\chi}^U, \varphi_n) = \sum_m \tilde{b}_m^U (1 - \tilde{N})_{mn}, \end{aligned}$$

where

$$\begin{aligned} N_{mn} &= (\phi_m^\dagger, \mathcal{P}_+ \otimes U^\dagger \delta U \varphi_n), \\ \tilde{N}_{mn} &= -(\phi_m^\dagger \mathcal{P}_- \otimes U^\dagger \delta U, \varphi_n). \end{aligned}$$

Therefore, the measure changes as $\prod da_n^U = \text{Det}_{V \otimes W_{\text{rt}} \otimes M} (1 - N) \prod da_n^{U+\delta U}$ and $\prod d\tilde{b}_n^U = \text{Det}_{V \otimes W_{\text{rt}} \otimes M} (1 - \tilde{N}) \prod d\tilde{b}_n^{U+\delta U}$, where M stands for the space for the eigenmodes of the Dirac operator. Noting that $\det(1 + \epsilon X) \sim e^{\epsilon \text{tr} X + O(\epsilon^2)}$ valid for small ϵ , we reach

$$\begin{aligned} \mathcal{D}\chi^U &= \mathcal{D}\chi^{U+\delta U} e^{-\delta\Gamma_\chi}, \\ \mathcal{D}\tilde{\chi}^U &= \mathcal{D}\tilde{\chi}^{U+\delta U} e^{-\delta\tilde{\Gamma}_\chi}, \end{aligned}$$

where $\delta\Gamma_\chi = \text{Tr}_{V \otimes W_{\text{rt}} \otimes M} N \equiv \int d^2x \sum_n \text{tr}_{V \otimes W_{\text{rt}}} N_{nn}$ and $\delta\tilde{\Gamma}_\chi = \text{Tr}_{V \otimes W_{\text{rt}} \otimes M} \tilde{N} \equiv \int d^2x \sum_n \text{tr}_{V \otimes W_{\text{rt}}} \tilde{N}_{nn}$. Here Tr_M stands for the trace for x and M . In principle, we can choose either Γ_χ or $\tilde{\Gamma}_\chi$ as the Jacobian in Eq. (4.6) if they can be regularized. In what follows, we will apply standard regularization scheme in Refs. [28,29]. Nevertheless, it turns out that they still give rise to divergent terms for each Γ . Fortunately, as we shall see below, they cancel each other in $\delta\Gamma_\chi + \delta\tilde{\Gamma}_\chi$. Therefore, it is suitable to define the Jacobian in Eq. (4.4) as a square root of the product of the naive Jacobians as $\text{Det}_{V \otimes W_{\text{rt}} \otimes M}^{\frac{1}{2}} (1 - N)(1 - \tilde{N})$. Namely,

$$\delta \ln J(U) \equiv \delta\Gamma = \frac{1}{2}(\delta\Gamma_\chi + \delta\tilde{\Gamma}_\chi).$$

It is possible to show that the Jacobian based on $\mathcal{D}\psi'\mathcal{D}\tilde{\psi}'$ for the Lagrangian (4.2) indeed yields the same Jacobian.

We now give the outline of how to calculate the Jacobian following Refs. [28,29]. Using the definition of the Tr_M , we have

$$\begin{aligned}
\delta\Gamma &= \frac{1}{2} \text{Tr}_{V \otimes W_{\text{rt}}} \sum_n \phi_n^\dagger(x) \mathcal{P} \otimes U^\dagger \delta U \varphi_n(x) \\
&= \frac{1}{2} \text{Tr}_{V \otimes W_{\text{rt}}} \sum_n \phi_n^\dagger(x) \mathcal{P} \otimes U^\dagger \delta U e^{-(\lambda_n/\Lambda)^2} \varphi_n(x) \\
&\equiv \frac{1}{2} \text{Tr}_{V \otimes W_{\text{rt}} \otimes M} \mathcal{P} \otimes U^\dagger \delta U e^{-(i\hat{P}/\Lambda)^2}
\end{aligned}$$

where ultraviolet cut-off Λ is introduced, which will be set $\Lambda \rightarrow \infty$ at the end of the calculation. Using the relation $\hat{P}^2 = \mathcal{P}_+ \hat{P}_- + \hat{P}_- \mathcal{P}_+$, we have

$$\begin{aligned}
\delta\Gamma &= \frac{1}{2} \text{Tr}_{V \otimes W_{\text{rt}} \otimes M} U^\dagger \delta U \left[\mathcal{P} (e^{\hat{P}\mathcal{P}/\Lambda^2} + e^{\mathcal{P}\hat{P}/\Lambda^2}) + (e^{\hat{P}\mathcal{P}/\Lambda^2} - e^{\mathcal{P}\hat{P}/\Lambda^2}) \right] \\
&= \frac{1}{2} \int \frac{d^2k}{(2\pi)^2} \text{Tr}_{V \otimes W_{\text{rt}}} e^{-ikx} U^\dagger \delta U \left[\mathcal{P} (e^{\hat{P}\mathcal{P}/\Lambda^2} + e^{\mathcal{P}\hat{P}/\Lambda^2}) + (e^{\hat{P}\mathcal{P}/\Lambda^2} - e^{\mathcal{P}\hat{P}/\Lambda^2}) \right] e^{ikx},
\end{aligned}$$

where in the last line we have switched the basis for M from the eigenmodes of the Dirac operator into plane waves. Expanding Λ and taking the limit $\Lambda \rightarrow \infty$, we reach

$$\delta\Gamma = \frac{1}{2} \int \frac{d^2k}{(2\pi)^2} e^{-k^2} \text{Tr}_{V \otimes W_{\text{rt}}} [\mathcal{P} \otimes U^\dagger \delta U (i\partial \times D) + 1 \otimes U^\dagger \delta U (\partial \cdot D)]. \quad (4.7)$$

where $a \times b = \sigma_{\mu\nu} a_\mu b_\nu$ and $a \cdot b = g_{\mu\nu} a_\mu b_\nu$ with

$$\sigma_{\mu\nu} = \frac{1}{2i} [\gamma_\mu, \gamma_\nu] = \epsilon_{\mu\nu} \sigma_3 \otimes \sigma_3, \quad g_{\mu\nu} = \frac{1}{2} \{\gamma_\mu, \gamma_\nu\} = \delta_{\mu\nu} 1 \otimes 1. \quad (4.8)$$

Therefore, taking trace in x - and V -space, we have

$$\delta\Gamma = \frac{1}{2\pi} \int d^2x \text{tr}_{W_{\text{rt}}} U^\dagger \delta U (\delta_{\mu\nu} - i\epsilon_{\mu\nu}) \partial_\mu A_\nu.$$

It is nowadays well-known that the action function satisfying above includes not only the usual kinetic term but also the WZW term,

$$\Gamma = S_{\text{WZW}}^k \equiv \frac{1}{4\pi} \int d^2x \text{tr}_{W_{\text{rt}}} \partial_\mu U \partial_\mu U^\dagger + \frac{ik}{12\pi} \int d^3x \epsilon_{\mu\nu\lambda} \text{tr}_{W_{\text{rt}}} U^\dagger \partial_\mu U U^\dagger \partial_\nu U U^\dagger \partial_\lambda U, \quad (4.9)$$

with $k = 2$. Adding the symmetry-breaking term in (4.6), we end up with an action

$$S = S_{\text{WZW}}^{k=2} + \frac{\epsilon m}{2g'} \int d^2x \text{tr}_{W_{\text{rt}}} (U + U^\dagger). \quad (4.10)$$

Therefore, it turns out that the Goldstone mode of the model with chiral symmetry of type $\mathcal{P}_1$ can be described by $\text{Sp}(n)_2$ WZW model (4.9) with a symmetry breaking term as in (4.10).

So far we have considered the case where Hamiltonian has the chiral symmetry specified by $\mathcal{P}_1$ in Eq. (2.7). As to the other cases, similar calculations lead to $\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model, which is so-called chiral orthogonal class or $B\text{DI}$, the same universality class as HWKM model.

V. PHASE DIAGRAM

We present a schematic phase diagram at the zero energy in Fig. 2, which is valid for weak disorder, i.e, for small g_j . In the case $g_3 = g_4 = 0$, which is denoted by a white surface in the figure, the model belongs to class CI and is described by $\text{Sp}(n)_2$ WZW model with some irrelevant perturbations discussed in Sec. IIIC and IIID. The reason why the present WZW model is level-2 is that the continuum limit (2.2) leads to double Dirac fermions. One of the main origin of the irrelevant interactions is that the lattice model has only on-site potentials without any hopping randomness. Namely, the model does not have maximum entropy. Another origin is the anisotropy $g_1 \neq g_2$, as discussed in Sec. IIID. Besides, the continuum model should have basically many higher order irrelevant interactions. Therefore, at the zero energy any points on this surface flow to the $\text{Sp}(n)_2$ WZW fixed point, denoted by a black




FIG. 2. Schematic phase diagram at the zero energy in the case of weak disorder. We omitted the axis g_4 . Black circle indicates the $\text{Sp}(n)_2$ WZW fixed point. Shaded surfaces stand for “critical surfaces” described by $\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model. Any points away from these three surfaces flow to strong coupling limit of $\text{O}(2n, 2n)/\text{O}(2n) \times \text{O}(2n)$ nonlinear sigma model.

circle in the figure. It turns out that around the fixed point the density of states obeys the scaling law $\rho(E) \sim E^{\frac{1}{4}}$, following from the exact results obtained by Knizhnik and Zamolodchikov [30].

On the other hand, the universality class of the two surfaces $g_1 = g_4 = 0$ and $g_2 = g_4 = 0$, shaded in the figure, is different. They belong to the class of $\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model or in other words, chiral orthogonal class or class $B\text{DI}$, which is the same universality class as that of HWKM model. It is well-known that the beta function of the renormalization group vanishes and the coupling constant of the nonlinear sigma model is not renormalized. Hence any points on these surfaces are critical. The density of states is conjectured to be divergent at the zero energy, according to $\rho(E) \sim \frac{\Lambda}{E} \exp[-\alpha \sqrt{\ln(\Lambda/E)}]$ with a positive constant α [1].

These surfaces, one CI and two $B\text{DI}$ ’s, are unstable: Out the surfaces renormalization group flows are toward the strong-coupling limit of $\text{Sp}(2n)/\text{Sp}(n) \times \text{Sp}(n)$ nonlinear sigma model, as shown by Fisher and Fradkin for $g_1 = g_2 = g_3 = g_4$ case. Therefore, all states away from the surfaces are expected to be localized.

VI. SUMMARY AND CONCLUDING REMARKS

We have studied the phase diagram of the disordered fermion model with random on-site potential defined on a two-dimensional square lattice with π -flux. The continuum limit near the zero energy yields Dirac fermions with random potentials specified by four independent coupling constants. It has been argued that in addition to time-reversal symmetry, this model has so-called chiral symmetry in three subspaces in the space of the coupling constants. Here, chiral symmetry means that there exist a matrix anticommuting with the Hamiltonian. What is interesting is that this model allows three kinds of chiral symmetry, which is due to the species doubling of the lattice fermion. It has been shown that one of them leads to $\text{Sp}(n) \times \text{Sp}(n)$ symmetry of the Lagrangian for generating functional of ensemble-averaged Green functions whereas the others lead to $\text{U}(2n)$ symmetry, both of which is broken to $\text{Sp}(n)$. It has been found that the fluctuation around the saddle point is described, respectively, by $\text{Sp}(n)_2$ WZW model for the former and $\text{U}(2n)/\text{Sp}(n)$ nonlinear sigma model for the latter. Based on these results, we have proposed a phase diagram.

The Dirac fermion model with $\mathcal{P}_1$ symmetry calculated in detail in the text is quite similar to that for the disordered d -wave superconductors with spin-rotation and time-reversal symmetries. Actually, effective Lagrangians for these systems have $\text{Sp}(n) \times \text{Sp}(n)$ symmetry and hence both the models belong to the same universality class CI . However, fixed-points are quite different due to the WZW term. It was shown [17,23] that in the d -wave superconductors the WZW terms for several species of Dirac fermions cancel each other and the resultant theory predicts localization of

quasiparticles at the band center. Contrary to this, the model treated in this paper is described by level-2 WZW model, which means that it has delocalized states. The difference lies in Eq. (4.8): In the present model the WZW term survives owing to the relation $\sigma_{\mu\nu} = [\gamma_\mu, \gamma_\nu]/2i = \epsilon_{\mu\nu}\mathcal{P}_1$, which is due to $\mathcal{P}_1 = i\gamma_1\gamma_2$. This is just the same definition of so-called γ_5 in the particle physics literature, although the present γ matrices are not 2×2 but 4×4 . Conversely, we can say that chiral symmetry only in this sense yields the chiral anomaly and hence the WZW term, although we often use the same words for other similar symmetry in the Anderson localization literature. Considering the fact that the lattice model having $\mathcal{P}_1$ chiral symmetry is realized in the case $v'_3 = -v'_1$ and $v'_4 = -v'_2$ in Eq. (2.1), which is indeed a fine-tuned model, we conclude that the delocalization due to the existence of the WZW term seldom occurs in nature.

* Email: fukui@mito.ipc.ibaraki.ac.jp

- [1] R. Gade, Nucl. Phys. **B398**, 499 (1993); R. Gade and F. Wegner, *ibid.*, **B360**, 213 (1991).
- [2] M. R. Zirnbauer, J. Math. Phys. **37**, 4986 (1996).
- [3] A. Altland and M. R. Zirnbauer, Phys. Rev. **B55**, 1142 (1997).
- [4] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
- [5] A. Furusaki, Phys. Rev. Lett. **82**, 604 (1999), and references therein.
- [6] A. Altland and B. D. Simons, Nucl. Phys. **B562**, 445 (1999).
- [7] S. Guruswamy, A. LeClair, and A. W. W. Ludwig, cond-mat/9909143.
- [8] M. Fabrizio and C. Castellani, Nucl. Phys. **B583**, 542 (2000).
- [9] Y. Hatsugai, X.-G. Wen, and M. Kohmoto, Phys. Rev. **B56**, 1061 (1997); Y. Morita and Y. Hatsugai, Phys. Rev. Lett. **79**, 3728 (1997); Y. Morita and Y. Hatsugai, Phys. Rev. **B58**, 6680 (1998).
- [10] T. Fukui, Nucl. Phys. **B562**, 477 (1999).
- [11] A. A. Nersisyan, A. M. Tsvelik, and F. Wenger, Phys. Rev. Lett. **72**, 2628 (1994).
- [12] K. Ziegler, M. H. Hettler, and P. J. Hirschfeld, Phys. Rev. Lett. **77**, 3013 (1996).
- [13] T. Senthil, M. P. A. Fisher, L. Balents, and C. Nayak, Phys. Rev. Lett. **81**, 4704 (1998); T. Senthil and M. P. A. Fisher, Phys. Rev. **B60**, 6893 (1999).
- [14] R. Bundschuh, C. Cassanello, D. Serban, and M. R. Zirnbauer, Phys. Rev. **B59**, 4382 (1999).
- [15] T. Senthil and M. P. A. Fisher, Phys. Rev. **B61**, 9690 (2000).
- [16] M. Bocquet, D. Serban, and M. R. Zirnbauer, cond-mat/9910480.
- [17] T. Fukui, cond-mat/0002002.
- [18] S. Vishveshwara and M. P. A. Fisher, cond-mat/0003018.
- [19] P. Fendley and R. M. Konik, cond-mat/0003436.
- [20] W. A. Atkinson, P. J. Hirschfeld, and A. H. MacDonald, cond-mat/0002333.
- [21] J.-X. Zhu, D. N. Sheng, and C. S. Ting, cond-mat/0005266.
- [22] W. A. Atkinson, P. J. Hirschfeld, A. H. MacDonald, and K. Ziegler, cond-mat/0005487.
- [23] A. Altland, B. D. Simons, and M. R. Zirnbauer, cond-mat/0006362.
- [24] A. M. M. Pruisken, Nucl. Phys. **B235**, 277 (1984); H. Levine, S. B. Libby, and A. M. M. Pruisken, *ibid* **B235**, 30, 49, 71 (1984).
- [25] M. P. A. Fisher and E. Fradkin, Nucl. Phys. **B251**, 457 (1985).
- [26] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, Phys. Rev. **B50**, 7526 (1994).
- [27] R. Bundschuh, C. Cassanello, D. Serban, and M. R. Zirnbauer, Nucl. Phys. **B532**, 689 (1998).
- [28] K. Fujikawa, Phys. Rev. **D21**, 2848 (1980); *ibid* **D22**, 1499 (1980) (E)
- [29] L. Alvarez-Gaumé and P. Ginsparg, Nucl. Phys. **B243**, 449 (1984).
- [30] V. G. Knizhnik and A. B. Zamolodchikov, Nucl. Phys. **B247**, 83 (1984).